

SF00-SF06-0006

SPARTAN

Spray On Sheep Blowfly Protection





Active Content & Action

Cyromazine 60 g/L. Cyromazine is part of the triazine group of chemicals. Cyromazine disrupts nerve transmission in the target parasite.

Use

A ready to use solution to protect sheep against fly strike by Lucilia Cuprina.

Indications

Nil

Period of protection

When applied as directed will protect long wool sheep against fly strike for 11 weeks.

Dosage and Administration External Use Only.

Do not dilute. Do not mix with any other product.

Except for small lambs under 15Kg, the dose volume is determined by the number of months shearing. Apply Spartan according to the diagrams and tables in the Directions for Use.

Spartan Spray may be applied to all breeds of sheep, providing protection against flystrike. Spartan Spray is applied as a low-volume treatment onto or near areas where protection is required. When applying, the nozzle of the applicator gun should be approximately 25cm above the fleece, to achieve a 15cm wide band at each pass.

Months Since Shearing	Total Volume/ Sheep (MI)	Volume Per Treatment Band (MI)	No. of Bands	Sheep Treated Per 1L	Critical Comments
For the prevention of body strike				1. Use a ZAGRO AUSTRALIA	
Up to 6	34	17	2	29	recommended Spray-On applicator gun. 2. Ensure the applicator gun is clean. 3. Calibate applicator gun using an accurate volume measure. 4. Ensure all areas to be protected are treated. 5. When preventing body strike ensure
7	38	19	2	26	
8	44	22	2	22	
9	50	25	2	20	
10	56	28	2	17	
Lambs less than 15kg	17	1 <i>7</i>	1	58	
For the prevention of body and crutch strike				bands overlap along	
Up to 6	51	1 <i>7</i>	3	19	the midline. 6. This product is not ideal for treating existing strikes. Treated larvae may take 3-4 days to die. 7. Heavy rain following the application of this product
7	57	19	3	17	
8	66	22	3	15	
9	75	25	3	13	
10	84	28	3	11	
Lambs less than 15kg	34	1 <i>7</i>	2	29	
FOR THE PREVENTION OF POLL STRIKE Rams and sheep susceptible to poll strike should be treated. Apply at least 5 mL of SPARTAN SPRAY-ON to the areas requiring protection, around the poll and horns.				could diminish the period of protection. 8. SPARTAN SPRAY-ON is coloured for band identification. This scourable dye will disappear shortly after treatment.	

Doses

400 doses per 20L pack (at 9 months shearing for body strike).

Pack Size(s) Available 20L

Australian Poisons Schedule S5

Presentation

440 drums to a pallet

APVMA Approval No.: 80858



In Agrisolutions Since 1953



Withholding Periods

Meat : 7 Days

Milk : Dispose of all milk collected from animals treated with this product

Wool : 2 months

Trade Advice

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 28 days before slaughter for export. Before using this product, confirm the current ESI from ZAGRO AUSTRALIA Pty Ltd on 1800 144 884 or the APVMA website (www.apvma.gov.au/residues)

Accessories

This product can be used with any spray on applicator designed to apply spray on products to sheep. Contact your local reseller for the different options.